

## LA-UR-19-28184

Approved for public release; distribution is unlimited.

Title: An Introduction to Quantum Dots as Scintillating  $\gamma$ -Radiation Detectors

Author(s): Starvaggi, Nicholas Charles

Intended for: Report

Issued: 2019-08-13

---

**Disclaimer:**

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

# An Introduction to Quantum Dots as Scintillating $\gamma$ -Radiation Detectors




**Nicholas Charles Starvaggi**  
**NEN-3: International Threat Reduction**  
Summer 2019 SULI Internship



Managed by Triad National Security, LLC for the U.S. Department of Energy's NNSA

# Scintillation Technology

- A **scintillator** is any material capable of converting some of the energy of incoming incident particles into low-energy photons in the ultraviolet to visible range for a variety of technical applications.<sup>1</sup>
  - Nuclear engineering
  - Medical imaging
  - National Security
  - High-energy physics
- Scintillating Radiation Detectors
  - Inorganic or organic composition
  - Luminescent properties
    - Identify radioactive sources
    - Characterize radiation
  - Quantify radiation intensity




**Fig. 1:** Mechanism for  $\gamma$ -radiation down-conversion with plastic scintillators.<sup>1</sup>

# Types of Scintillators

- Inorganic Crystals<sup>1</sup>
  - NaI(Tl), CsI(Na)
    - High light yields for  $\gamma$ -radiation detection and spectroscopy
    - Expensive and susceptible to rapid degradation
- Organic Materials<sup>1</sup>
  - Crystals
    - Acceptable light yields
    - Fragile, difficult to machine and polish
  - Liquids
    - Large-scale production possible
    - Flammability and toxicity of aromatic solvents
  - Plastics
    - Low light yields
    - Minimal hazards



**Fig. 2:** NaI(Tl) scintillator. Courtesy of [www.ost-photonics.com](http://www.ost-photonics.com)



**Fig. 3:** Luminescent plastic scintillators. Courtesy of [www.oaaw.ac](http://www.oaaw.ac)

# Nal(Tl) Scintillator Schematic





Fig. 4: NaI(Tl) Scintillator and PMT schematic. Courtesy of [www.nuclear-power.net](http://www.nuclear-power.net)

# Scintillating $\gamma$ -Radiation Detectors

- Ideal Scintillator Characteristics<sup>2</sup>
  - High-Z composition
    - Increase  $\gamma$ -attenuation
    - Generate photoelectron production
  - High light yield
    - Increased photon count per unit of energy deposited
    - Optical clarity
  - Short emission decay lifetime
    - Timing resolution<sup>4</sup>
    - Immediate system response
  - Low Cost
  - Ease of large-scale fabrication




**Fig. 5:** Effect of organo-bismuth (high-Z) loading on the radiation response of a scintillator.<sup>3</sup>

# Wavelength Shifters for Advanced Scintillators



**Fig. 6:** Mechanism for  $\gamma$ -radiation down-conversion with fluor-doped plastic scintillators.<sup>1</sup>




**Fig. 7:** Typical responsivity (A/W) of an avalanche photodiode (APD). Courtesy of [www.thorlabs.com](http://www.thorlabs.com).

Quantum dots offer an alternate method to  $\gamma$ -radiation downconversion in the next generation of photomultiplication devices.



# QD Basics


- Semiconductor Nanoparticle Crystals
  - Inorganic Core
    - High-Z element composition
    - Interactions with radiation
  - Ligand Shell
    - Easily manipulated
    - Determines solubility<sup>5</sup>
- Quantum Mechanical Properties
  - Particle size
    - 2-10 nm
  - Energy bandgap
  - Emission profile<sup>6</sup>




*Fig. 8: Ball-and-stick model of a QD showing the inorganic core and ligand shell. Fig. adapted from proposal.<sup>6</sup>*

# Key Properties

- Characteristic Emission Color
  - Dependent on particle size and bandgap
- Smaller QDs, Larger Bandgap
  - More energy to promote a single  $e^-$  to an excited state
  - Results in a higher energy frequency ( $\nu$ ) and shorter wavelength ( $\lambda$ )
  - Emit light towards violet/indigo end of the visible light spectrum
- Larger QDs, Smaller Bandgap
  - Less energy to promote a single  $e^-$  to an excited state
  - Results in lower  $\nu$  and longer  $\lambda$
  - Emit light towards red/orange end of the visible light spectrum




**Fig. 9:** Comparison of the energy orbital diagrams of molecules, QDs, and semi-conductors.




**Fig. 10:** QD sizes displayed along the visible light spectrum.<sup>6</sup>

# Synthesis of QDs

- Basic Procedure
  - Pyrolysis (decomposition) of organometallic precursors in hot solvent generates QDs<sup>5</sup>
    - QDs are bound in aqueous solution
    - Subsequent reactions in coordinating solvents are often performed to manipulate the ligands
  - Reaction length affects nanoparticle size
- Special Focus on High-Z Element Combinations
  - Cd/Se, Cd/S
  - Pb/Se, PbS
  - In/As, In/P
  - Zn/Se, Zn/S




**Fig. 11:** Effect of reaction time in the synthesis of QDs.<sup>6</sup>



**Fig. 12:** Diagram of a PbSe/CdSe core-shell quantum dot.<sup>7</sup>

# QDs as Scintillating $\gamma$ -Radiation Detectors


- QD Research
  - Nanometer dimensions increases bandgap and allows for visible luminescence<sup>9</sup>
  - Integrates necessary high-Z inorganics with organic dyes in a polymer matrix<sup>8</sup>
    - Polystyrene (PS) and polyvinyl toluene (PVT)
    - Potential copolymerization applications with acrylate compounds to increase durability
  - Resulting plastic scintillator is low-cost and has a short-emission decay lifetime
    - Optical transparency contributes to high light yield
- Challenges
  - Self-attenuation (self-reabsorption)
  - Incorporation into a polymer matrix



*Fig. 13: Self-attenuation in average quantum dots. Fig. adapted from proposal.<sup>6</sup>*

# Giant Quantum Dots (gQDs) in LANL Scintillation Technology


- gQDs, developed by Jennifer Hollingsworth at CINT, are a LANL-patented alternative to standard QDs<sup>10</sup>
  - Ultra-thick CdSe/CdS core-shell system
  - Increased photostability
- Special Core-Shell Functions
  - Shell absorbs light, core emits light
  - Absorption energy is higher than emission energy
    - Shell serves as the wider-gap semiconductor in comparison to the core material
  - Light emitted by a single gQD is thus not reabsorbed by other gQDs in polymer matrix



*Fig. 14: Comparison of standard QDs (left) and LANL-patented gQDs (right). The absorption (dark lines) and emission (light lines) demonstrate the lack of self-attenuation in gQDs. Fig. adapted from proposal.<sup>6</sup>*


# Research Goals, Challenges, & Solutions

- Polymerization of PS and PVT
  - Initial photoinitiation via BAPO
  - Copolymerization with 2-carboxyethyl acrylate ( $C_6H_8O_4$ )
    - 3:1 Ratio
    - 6:1 Ratio
- Incorporate wavelength shifters
  - POPOP ( $C_{24}H_{16}N_2O_2$ )
  - Characterization
    - IR Spectroscopy
    - UV-VIS spectrophotometry
    - Fluorimetry




**Fig. 15:** 2CEA:PS/PVT (6:1) samples with 3 mg of POPOP exposed to UV light.

# Initial IR Results



**Fig. 16:** IR spectra of 2CEA/PS composite (black line) and neat styrene (orange line). Styrene IR courtesy of [www.webbook.nist.gov](http://www.webbook.nist.gov).

# IR Results (cont.)



Group	Range ( $\text{cm}^{-1}$ )
$\text{R}-\text{CH}=\text{CH}_2$	985-1000 and 905-920
$\text{R}_2\text{CH}=\text{CH}_2$	880-900


**Table 1:** IR frequency ranges ( $\text{cm}^{-1}$ ) for functional groups relevant to styrene.

**Fig. 17:** Zoomed-in insert of Fig. 15 from 500-1000  $\text{cm}^{-1}$ . Styrene IR courtesy of [www.webbook.nist.gov](http://www.webbook.nist.gov).



# Conclusions & Future Work

- Inconclusive polymerization trials
  - IR data suggests a lack of polymerization
  - Extended exposure to visible light resulted in solid composite
  - Vinyl toluene and styrene performed identically
- Future Work
  - Cationic photopolymerization
  - Incorporate gQDs to mitigate self-attenuation
    - UV-VIS Spectrophotometry
    - Fluorimetry
  - Stronger UV exposure to aid photopolymerization



*Fig. 18: Solid styrene/BAPO composite after seven days of visible light exposure.*

# Acknowledgements

- Special Thanks To...
  - Dr. Amanda Graff
  - Dr. Morag Smith
  - NEN-3: International Threat Reduction
  - NEN-1: Safeguards Science & Technology
  - MST-7: Engineered Materials
  - Workforce Development for Teachers and Scientists (WDTs)
  - Science Undergraduate Laboratory Internships (SULI) Program
  - Los Alamos National Laboratory (LANL)



U.S. DEPARTMENT OF  
**ENERGY**

Office of  
Science



# Questions?

# References

- <sup>1</sup>Hajagos, T. J., Liu, C., Cherepy, N. J., Pei, Q., *Adv. Mater.* **2018**, 30, 1706956.
- <sup>2</sup>Chao Liu, Zhou Li, Tibor Jacob Hajagos, David Kishpaugh, Dustin Yuan Chen, and Qibing Pei. *ACS Nano* **2017** 11 (6), 6422-6430
- <sup>3</sup>Cherepy, N. J.; Sanner, R. D.; Beck, P. R.; Swanberb, E. L.; Tillotson, T. M.; Payne, S. A.; Hurlbut, C. R. *Nucl. Instrum. Methods Phys. Res., Sect. A* **2015**, 778, 126.
- <sup>4</sup>Yanagida, T. Inorganic scintillating materials and scintillation detectors. *Proc. Jpn. Acad., Ser. B* **2018**, 94, (2), 75-97.
- <sup>5</sup>Hollingsworth, J. A. *Encycl. Inorg. Bioinorg. Chem.* **2011**, 1.
- <sup>6</sup>Zhao, H.; Rosei, F. *Chem* **2017**, 3, 229.
- <sup>7</sup>Torchynska, T., Vorobiev, Y. Semiconductor II-VI Quantum Dots with Interface States and their Biomedical Applications. *Adv. Bio. Eng.* **2011**.
- <sup>8</sup>Murphy, C. J., & Coffer, J. L. (2002). Quantum Dots: A Primer. *Applied Spectroscopy*, 56(1), 16A-27A.
- <sup>9</sup>S. E. Létant\* and T.-F. Wang. *Nano Letters* **2006** 6 (12), 2877-2880
- <sup>10</sup>Yagnaseni Ghosh, Benjamin D. Mangum, Joanna L. Casson, Darrick J. Williams, Han Htoon, and Jennifer A. Hollingsworth. *Journal of the American Chemical Society* **2012** 134 (23), 9634-9643